

Immunise – their best protection

Talk to your doctor or practice nurse.



Pregnancy

1 injection, between 28-38 weeks covers:

tetanus
diphtheria
whooping cough (pertussis)

1 injection covers:

influenza



6

Weeks

1 dose by mouth covers:

rotavirus

1 injection covers:

diphtheria
tetanus
whooping cough (pertussis)
polio
hepatitis B
Haemophilus influenzae type b (Hib)

1 injection covers:

pneumococcal disease



3

Months

1 dose by mouth covers:

rotavirus

1 injection covers:

diphtheria
tetanus
whooping cough (pertussis)
polio
hepatitis B
Haemophilus influenzae type b (Hib)

1 injection covers:

pneumococcal disease



5

Months

1 injection covers:

diphtheria
tetanus
whooping cough (pertussis)
polio
hepatitis B
Haemophilus influenzae type b (Hib)

1 injection covers:

pneumococcal disease



15

Months

1 injection covers:

Haemophilus influenzae type b (Hib)

1 injection covers:

measles
mumps
rubella

1 injection covers:

pneumococcal disease

1 injection covers:

chickenpox (varicella)



4

Years

1 injection covers:

diphtheria
tetanus
whooping cough (pertussis)
polio

1 injection covers:

measles
mumps
rubella



11 & 12
Years

1 injection covers:

tetanus
diphtheria
whooping cough (pertussis)

2 injections, 6 months apart covers*:

human papillomavirus (HPV)

It's free. Immunise on time, every time. For more information visit www.health.govt.nz/immunisation

*Those aged 15 years or older need 3 doses.

This resource is available from health.govt.nz or the Authorised Provider at your local DHB. Revised March 2017. Code HE1221